

Oklahoma OSTP Grade 8 Math Summer Workbook

8-Week Skills Practice with Quick Reviews and Answers

Dr. A. Nazari

Copyright © 2026 Dr. A. Nazari

PUBLISHED BY VIEW MATH EDUCATION

VIEWMATH.COM

All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the author, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law, including Section 107 or 108 of the 1976 United States Copyright Act.

The information in this book is distributed on an “as is” basis, without warranty. While every precaution has been taken in the preparation of this work, neither the author nor the publisher shall have any liability to any person or entity with respect to any loss or damage caused or alleged to be caused directly or indirectly by the information contained in this book.

Copyright © 2026

SUMMER PRACTICE

Grade 8 Math Summer Workbook

Quick Reviews, Workbook Practice, Mixed Reviews, and Answers

This workbook keeps Grade 8 math fresh with short review lessons and strong daily practice. Each week moves from focused skills to a Friday mixed review, so students can remember the method, choose the right strategy, and explain their thinking.

- ✔ Read the Quick Review.
- ✎ Complete the workbook practice.
- ✔ Use Friday for mixed review.
- 💡 Study the answer explanations after trying first.

How to Use This Workbook

Use one short lesson at a time, then check and learn from mistakes.

The Weekly Routine

- Monday-Thursday** Read the Quick Review, then complete one workbook practice set.
- Friday** Complete the weekly mixed review to connect the week's skills.
- After practice** Check answers and read each explanation for any problem that felt uncertain.

For students

Try the problems before looking at the answer key. If you miss a problem, read the explanation, correct your work, and mark the skill to revisit later.

For parents and teachers

The workbook pages are designed for steady practice. Use the Quick Review for a short reteach, and use the answer explanations as the teaching step after a mistake.

Best practice habit

Show your reasoning. Grade 8 math is easier to check when you label variables, mark points on a graph, write equations, and include units or vocabulary in your answer.

Goal

By the end of 8 weeks, students will have reviewed major Grade 8 skills with workbook-style practice, visual models, mixed reviews, and teaching answer explanations.

What's Inside?

An 8-week workbook plan for Grade 8 summer math practice.

Week 1

Rational and irrational numbers, roots, integer exponents, scientific notation.

Week 2

Square and cube roots, scientific notation operations, proportional graphs.

Week 3

Slope, equations of lines, one-variable equations, systems of equations.

Week 4

Real-world systems, functions, comparing functions, linear and nonlinear patterns.

Week 5

Linear functions, graph descriptions, rigid transformations, congruence.

Week 6

Coordinate transformations, similarity, dilations, angles, Pythagorean theorem.

Week 7

Distance, volume formulas, scatter plots, trend lines, data interpretation.

Week 8

Linear models, two-way tables, variability, probability, and cumulative review.

Quick Reviews

Each topic begins with a compact review of the ideas, formulas, and methods students need before starting the practice.

Weekly Reviews

Friday pages mix the week's skills so students practice choosing the right method instead of following only one pattern.

Workbook Practice

Practice sets include computation, word problems, graphs, tables, diagrams, models, and short reasoning.

Answers

The answer key includes explanations that show the method and reasoning, not just the final answer.

The goal is steady confidence, not rushed memorization.

PREVIEW



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)





Grade 8 Reference Sheet



High-yield formulas and reminders for quick review

Exponents

$$a^m \cdot a^n = a^{m+n}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

$$(a^m)^n = a^{mn}$$

$$(ab)^n = a^n b^n$$

$$a^0 = 1 \text{ for } a \neq 0$$

$$a^{-n} = \frac{1}{a^n}$$

Scientific Notation

$$a \times 10^n, \text{ where } 1 \leq |a| < 10$$

Multiply: multiply decimals, add exponents.

Divide: divide decimals, subtract exponents.

Positive exponent: number is large.

Negative exponent: number is small.

Roots and Real Numbers

Perfect squares: 1, 4, 9, 16, 25, 36

49, 64, 81, 100, 121, 144

Perfect cubes: 1, 8, 27, 64, 125

$$\sqrt{2} \approx 1.414, \sqrt{3} \approx 1.732, \pi \approx 3.14159$$

Irrational numbers are nonrepeating, nonterminating decimals.

mals.

Slope and Lines

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\text{rise}}{\text{run}}$$

Slope-intercept form: $y = mx + b$

m is slope, b is the y -intercept.

Proportional relationship: $y = mx$ and graph passes

through $(0, 0)$.

Systems of Equations

A solution is the point where two graphs meet.

One solution: different slopes.

No solution: same slope, different intercepts.

Infinitely many solutions: same line.

Use substitution or elimination to solve exactly.

Functions

Each input has exactly one output.

Use the vertical line test on graphs.

Linear functions have a constant rate of change.

In $y = mx + b$, the initial value is b .

Compare functions by rate of change and initial value.

Transformations

Translation: slide. Reflection: flip. Rotation: turn.

These rigid motions keep lengths and angle measures.

Dilation: resize by scale factor k .

$k > 1$ enlarges; $0 < k < 1$ reduces.

Congruent figures are same size and shape. Similar figures

have proportional sides.

Angle Relationships

Triangle angle sum: 180°

Exterior angle = sum of the two remote interior angles.

Vertical angles are congruent.

With parallel lines, alternate interior angles are congruent.

Same-side interior angles sum to 180° .

Pythagorean Theorem

For a right triangle, $a^2 + b^2 = c^2$.

c is the hypotenuse, the longest side.

Distance formula: $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Use the theorem to check whether a triangle is right.

Volume

$$\text{Cylinder: } V = \pi r^2 h$$

$$\text{Cone: } V = \frac{1}{3} \pi r^2 h$$

$$\text{Sphere: } V = \frac{4}{3} \pi r^3$$

Use the same cubic unit as the dimensions.

If dimensions scale by k , volume scales by k^3 .

Scatter Plots

Positive association: points trend upward.

Negative association: points trend downward.

No association: no clear pattern.

A line of fit models the trend.

Slope describes the average rate of change.

Two-Way Tables

Rows and columns group categorical data.

Relative frequency = part divided by total.

Use row, column, or table totals depending on the question.

Compare percentages, not just counts.

Variability and Probability

Mean absolute deviation measures typical distance from the

mean.

$$\text{MAD} = \frac{\text{sum of absolute deviations}}{\text{number of values}}$$

Probability: $0 \leq P(\text{event}) \leq 1$

Experimental probability uses results. Theoretical probability

uses equally likely outcomes.

Table of Contents

8-week review plan

Week 1	The Number System	1
Week 2	Proportional Graphs	8
Week 6	Geometry and the Pythagorean Theorem	12
	Answer Key & Explanations	15

WEEK

1

The Number System

This Week's Days

Week 1 Day 1: Rational and Irrational Numbers	2
Week 1 Day 2: Estimating Irrational Numbers	4
Week 1 Day 5: Week 1 Mixed Review	6



Day 1 Rational and Irrational Numbers

SKILL SNAPSHOT

Real numbers include rational and irrational numbers. Rational numbers can be written as a ratio of integers; irrational numbers cannot.

- ✓ Natural numbers are counting numbers; whole numbers also include 0.
- ✓ Integers include positive numbers, negative numbers, and 0.
- ✓ Fractions, terminating decimals, and repeating decimals are rational.
- ✓ Square roots of perfect squares are rational; square roots of non-perfect squares are irrational.
- ✓ A repeating decimal can be converted to a fraction by multiplying, subtracting, and solving.

Remember: Classify by value, not by appearance. For example, $\sqrt{36}$ is rational because it equals 6.

☰ Classify real numbers.

- 1 Classify $\sqrt{64}$ as rational or irrational.

- 2 Classify $\sqrt{28}$ as rational or irrational.

- 3 Is -11 natural, whole, integer, rational, or irrational? List all that apply.
- 4 What is the most specific set for 0?
- 5 Which number is irrational?
 A. $\frac{5}{8}$ B. $0.\overline{27}$
 C. $\sqrt{45}$ D. $\sqrt{49}$
- 6 True or False: Every integer is rational.
 True False

☰ Use a real-number hierarchy.

- 7 Use the chart to place each number in its most specific listed set: 9, 0, -4 , $\frac{3}{5}$, $\sqrt{10}$.

Natural	Whole	Integer	Rational	Irrational
_____	_____	_____	_____	_____

- 8 Complete the statement: $\text{Natural} \subset \text{Whole} \subset \text{_____} \subset \text{Rational}$.
- 9 A student says $\sqrt{81}$ is irrational because it has a radical sign. Explain the error.



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



10 Which list contains only rational numbers?

A. $\sqrt{2}$, 4, 0.6

B. -7 , $\frac{1}{3}$, 2.5

C. π , 0, $\sqrt{16}$

D. $\sqrt{5}$, 9, $0.\bar{1}$

 **Convert repeating decimals.**

11 Convert $0.\bar{5}$ to a fraction. _____

13 Convert $0.4\bar{2}$ to a fraction. _____

12 Convert $0.\bar{1}8$ to a fraction. _____

14 True or False: $0.\bar{9} = 1$. True False



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



Day 2 Estimating Irrational Numbers

SKILL SNAPSHOT

Estimate square roots by trapping the number between nearby perfect squares, then deciding where it belongs on a number line.

- ✓ If $a^2 < n < b^2$, then \sqrt{n} is between a and b .
- ✓ Use perfect-square landmarks such as 36, 49, 64, 81, and 100.
- ✓ For nearest tenth estimates, square tenths such as 7.2^2 or 7.3^2 .
- ✓ Compare \sqrt{n} to a decimal by squaring the decimal.
- ✓ A square-root estimate should be near the root, not near the radicand.



Remember: For example, $36 < 42 < 49$, so $6 < \sqrt{42} < 7$. Since 42 is closer to 36 than to 49, the root is closer to 6.

Find the whole-number bounds.

- 1 $\sqrt{27}$ is between which two consecutive whole numbers? _____
- 2 $\sqrt{90}$ is between which two consecutive whole numbers? _____
- 3 $\sqrt{8}$ is between which two consecutive whole numbers? _____
- 4 Use the chart to bound $\sqrt{55}$.

7^2	8^2	9^2	10^2
49	64	81	100

Estimate to the nearest whole number or tenth.

- 5 Estimate $\sqrt{35}$ to the nearest whole number.

- 6 Estimate $\sqrt{70}$ to the nearest whole number.

- 7 Estimate $\sqrt{14}$ to the nearest tenth.

- 8 Estimate $\sqrt{65}$ to the nearest tenth.

- 9 Estimate $\sqrt{115}$ to the nearest tenth.

- 10 Estimate $\sqrt{6}$ to the nearest tenth.



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



Use number lines and comparisons.

11 Use the number line. Which is the best estimate for $\sqrt{52}$? _____



12 Use the number line. Is $\sqrt{30}$ greater than or less than 5.5? _____



13 Fill in $<$, $>$, or $=$. $\sqrt{41}$ _____ 6.4

14 Order from least to greatest: 4.6, $\sqrt{20}$, 4.4. _____

Apply the estimate.

15 A square patio has area 95 ft^2 . Estimate its side length to the nearest tenth. _____

16 Which is the best estimate for $\sqrt{3}$?

A. 1.2


B. 1.7

C. 2.3

D. 3.0



Find more at [ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



Day 5 Week 1 Mixed Review

WEEKLY SKILL CHECK

This review connects real-number classification, square roots, irrational estimates, and exponent rules.


- ✓ Simplify a square root before deciding whether it is rational or irrational.
- ✓ Repeating decimals are rational because they can be written as fractions.
- ✓ Use nearby perfect squares to estimate irrational numbers.
- ✓ Add exponents when multiplying like bases, subtract when dividing, and multiply for powers of powers.
- ✓ A negative exponent means reciprocal, not a negative value.

Remember: For mixed questions, identify the rule first, then compute carefully.

Real numbers and roots.

- 1 Classify $\sqrt{121}$ as rational or irrational.

- 2 Classify $\sqrt{30}$ as rational or irrational.

- 3 Which number is the most specific whole number: -2 , 0 , $\frac{1}{2}$, $\sqrt{7}$? _____
- 4 Convert $0.\bar{3}$ to a fraction. _____
- 5 Use the number line to estimate $\sqrt{45}$ to the nearest whole number.

 $6^2 = 36$ 45 $7^2 = 49$
- 6 Order from least to greatest:
 $\sqrt{10}$, 3.5 , $\sqrt{16}$.

Exponent rules.

- 7 Simplify $a^3 \cdot a^8$. _____
- 8 Simplify $\frac{z^{12}}{z^5}$. _____
- 9 Simplify $(n^4)^3$. _____
- 10 Write 2^{-5} with a positive exponent. _____
- 11 Which expression equals 10^4 ?
A. $10^7/10^3$ **B.** $10^2 \cdot 10^3$
C. $(10^4)^2$ **D.** 10^{-4}
- 12 True or False: $8^0 = 0$. True False



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



 **Mixed applications.**

13 A square has area 75 square inches. Is its side length rational or irrational?

14 Use the table to choose the best estimate for $\sqrt{58}$.

7^2	58	8^2
49	58	64

A. 7.1

B. 7.6

C. 8.4

D. 9.0

15 Simplify $\frac{(x^3)^4}{x^5}$. _____

16 Convert $0.\overline{16}$ to a fraction. _____



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



WEEK

2

Proportional Graphs

 This Week's Days 

Week 2 Day 4: Graphing Proportional Relationships 9



Day 4 Graphing Proportional Relationships

SKILL SNAPSHOT

A proportional relationship has equation $y = kx$. Its graph is a straight line through the origin, and k is the constant of proportionality.

- ✓ In a table, divide y by x for each nonzero x .
- ✓ In an equation, proportional relationships have the form $y = kx$ with no added constant.
- ✓ On a graph, the line must pass through $(0, 0)$.
- ✓ The constant k is also the slope or unit rate.
- ✓ Use $y = kx$ to find missing values once k is known.



Remember: A straight line can be linear without being proportional. Proportional graphs must include the origin.

Find the constant of proportionality.

- 1 In a proportional relationship, $x = 4$ and $y = 18$. Find k . _____
- 2 A graph of a proportional relationship passes through $(6, 15)$. Find the unit rate. _____
- 3 Use the table to find k . _____

x	2	5	8
y	7	17.5	28

- 4 Use the table. Is the relationship proportional? _____

x	1	3	6
y	4	12	24

Interpret graphs.

- 5 Does the graph show a proportional relationship? _____



Get Online



Find more at ViewMath.com/Grade8





6 Does the graph show a proportional relationship? _____



7 Use the graph through $(0, 0)$ and $(5, 20)$. What is the equation? _____

☰ Equations and missing values.

8 Which equation is proportional?

A. $y = 7x$

B. $y = 7x + 2$

C. $y = x - 7$

D. $y = 7$

9 Is $y = 0.8x$ proportional? _____

10 Is $y = 3x - 5$ proportional? _____

11 If $y = 2.25x$, find y when $x = 12$.

☰ Applications and comparisons.

12 A printer produces 45 pages in 3 minutes at a constant rate. Write the proportional equation.

13 A cyclist travels according to $y = 14x$, where x is hours and y is miles. Another cyclist travels 45 miles in 3 hours. Who is faster? _____

14 True or False: A proportional relationship can include the point $(0, 5)$.

True

False

15 Use the table. Which store has the lower unit price? _____

Store	Pounds	Cost
A	4	\$18
B	6	\$24



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



- 16 A line through the origin contains $(9, 6)$. Find y when $x = 15$. _____

PREVIEW



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



WEEK

6

Geometry and the
Pythagorean Theorem

 This Week's Days 

Week 6 Day 4: The Pythagorean Theorem **13**



Day 4 The Pythagorean Theorem

SKILL SNAPSHOT

In a right triangle, the legs a and b and hypotenuse c satisfy $a^2 + b^2 = c^2$. The hypotenuse is opposite the right angle and is always the longest side.

- To find a hypotenuse, add the squares of the legs.
- To find a missing leg, subtract the known leg square from the hypotenuse square.
- The converse checks whether three side lengths form a right triangle.
- Round only at the end when a square root is not exact.



Find missing sides.

1 A right triangle has legs 9 and 12. Find the hypotenuse. _____

2 A right triangle has legs 7 and 24. Find the hypotenuse. _____

3 A right triangle has hypotenuse 13 and one leg 5. Find the other leg. _____

4 A right triangle has hypotenuse 20 and one leg 16. Find the other leg. _____

5 Find the missing side.



6 Find the missing leg when $c = 26$ and the other leg is 10. _____

Use the converse.

7 Do side lengths 10, 24, 26 form a right triangle? _____

8 Do side lengths 8, 14, 17 form a right triangle? _____

9 Which set forms a right triangle?

A. 5, 6, 7

B. 6, 8, 10

C. 7, 8, 9

D. 9, 10, 12



Get Online



Find more at ViewMath.com/Grade8



10 True or False: In a right triangle, the hypotenuse can be shorter than a leg.

True

False

11 A triangle has side lengths 9, 9, 13. Is it a right triangle? _____

 **Apply the theorem.**

12 A ladder is 15 ft long. Its foot is 9 ft from a wall. How high does it reach? _____

13 A rectangular field is 30 m by 40 m. Find the diagonal distance across the field. _____

14 A ramp rises 3 ft over a horizontal run of 11 ft. Find the ramp length to the nearest tenth.

 **Coordinate and 3D uses.**

15 Find the distance between (1, 2) and (7, 10).

16 A box is 6 cm by 8 cm by 24 cm. Find the space diagonal. _____



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



ANSWER KEY

Answer Key & Explanations

Use the answers to check your work, then read the explanations to learn the method.

✔ How to review

First compare your final answer. If it does not match, read the explanation slowly and redo the problem beside it. The goal is to understand the move that gets you from the question to the final answer.

Week 1 Day 1: Rational and Irrational Numbers

Answers

1 Rational

2 Irrational

3 Integer, rational, real

4 Whole number

5 C

6 True

7 9; 0; -4 ; $\frac{3}{5}$; $\sqrt{10}$

8 Integer

9 $\sqrt{81} = 9$, so it is rational

10 B

11 $\frac{5}{9}$ 12 $\frac{2}{11}$ 13 $\frac{19}{45}$

14 True

Explanations

1 The radical simplifies because 64 is a perfect square. Since $\sqrt{64} = 8$, the value can be written as $\frac{8}{1}$.

2 The number 28 is not a perfect square, so $\sqrt{28}$ does not simplify to a rational number. Its decimal form does not terminate or repeat.

3 The number -11 is an integer because it is a negative whole-number value. It is also rational because $-11 = \frac{-11}{1}$.

4 The number 0 is a whole number and also an integer and rational. It is not usually counted as a natural number in this course.

5 45 is not a perfect square, so $\sqrt{45}$ is irrational. The fraction, repeating decimal, and $\sqrt{49} = 7$ are rational.

6 Any integer n can be written as $\frac{n}{1}$. That form is a ratio of integers, so every integer is rational.

7 Choose the smallest set that fits each value. The square root of 10 is irrational because 10 is not a perfect square.

8 The nested order is natural, whole, integer, rational. Each set contains all numbers from the set before it.

9 The classification depends on the value after simplification. Since 81 is a perfect square, the radical equals the integer 9.



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



- 10 Integers, fractions, and terminating decimals are rational. The other lists include $\sqrt{2}$, π , or $\sqrt{5}$, which are irrational.
- 11 Let $x = 0.\overline{5}$. Then $10x = 5.\overline{5}$, so subtracting gives $9x = 5$ and $x = \frac{5}{9}$.
- 12 Let $x = 0.\overline{18}$. Then $100x = 18.\overline{18}$, so $99x = 18$ and $x = \frac{18}{99} = \frac{2}{11}$.
- 13 Let $x = 0.4222\dots$. Then $100x = 42.222\dots$ and $10x = 4.222\dots$, so $90x = 38$ and $x = \frac{19}{45}$.
- 14 Let $x = 0.\overline{9}$, so $10x = 9.\overline{9}$. Subtracting gives $9x = 9$, so $x = 1$.

📅 Week 1 Day 2: Estimating Irrational Numbers

✓ Answers

- 1 5 and 6 2 9 and 10 3 2 and 3 4 7 and 8 5 6 6 8 7 3.7 8 8.1
- 9 10.7 10 2.4 11 7.2 12 Less than 5.5 13 $>$ 14 4.4, $\sqrt{20}$, 4.6 15 9.7 ft
- 16 B

💡 Explanations

- 1 Use nearby perfect squares: $25 < 27 < 36$. Taking square roots gives $5 < \sqrt{27} < 6$.
- 2 $81 < 90 < 100$, so $\sqrt{90}$ is between $\sqrt{81} = 9$ and $\sqrt{100} = 10$.
- 3 $4 < 8 < 9$, so $\sqrt{8}$ is between 2 and 3. The root is closer to 3 because 8 is close to 9.
- 4 The chart shows $49 < 55 < 64$. Therefore $\sqrt{55}$ is between 7 and 8.
- 5 35 is between 25 and 36, and it is much closer to 36. Therefore $\sqrt{35}$ rounds to 6.



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



- 6 70 is between $64 = 8^2$ and $81 = 9^2$. It is closer to 64, so $\sqrt{70}$ rounds to 8.
- 7 $3.7^2 = 13.69$ and $3.8^2 = 14.44$. Since 14 is closer to 13.69, $\sqrt{14} \approx 3.7$.
- 8 $8.0^2 = 64$ and $8.1^2 = 65.61$. The value 65 is closer to 65.61 than to 64, so the nearest tenth estimate is 8.1.
- 9 $10.7^2 = 114.49$ and $10.8^2 = 116.64$. Since 115 is closer to 114.49, the nearest tenth estimate is 10.7.
- 10 $2.4^2 = 5.76$ and $2.5^2 = 6.25$. The value 6 is closer to 5.76, so $\sqrt{6} \approx 2.4$.
- 11 $7.2^2 = 51.84$, which is very close to 52. The number line places $\sqrt{52}$ just to the right of 7.2.
- 12 $5.5^2 = 30.25$, which is greater than 30. Therefore $\sqrt{30}$ must be slightly less than 5.5.
- 13 $6.4^2 = 40.96$, which is slightly less than 41. Since 6.4 squared is too small, $\sqrt{41}$ is slightly greater than 6.4.
- 14 $\sqrt{20} \approx 4.5$, because $4.5^2 = 20.25$. That places it between 4.4 and 4.6.
- 15 The side length is $\sqrt{95}$. Since $9.7^2 = 94.09$ and $9.8^2 = 96.04$, $\sqrt{95}$ is about 9.7 ft.
- 16 $\sqrt{3}$ is between 1 and 2. Since $1.7^2 = 2.89$ and $1.8^2 = 3.24$, the best choice is 1.7.

📅 Week 1 Day 5: Week 1 Mixed Review

✔ Answers

- 1 Rational 2 Irrational 3 0 4 $\frac{1}{3}$ 5 7 6 $\sqrt{10}$, 3.5, $\sqrt{16}$ 7 a^{11} 8 z^7
 9 n^{12} 10 $\frac{1}{32}$ 11 A 12 False 13 Irrational 14 B 15 x^7 16 $\frac{1}{6}$

💡 Explanations



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



- 1 121 is a perfect square, so $\sqrt{121} = 11$. Since $11 = \frac{11}{1}$, it is rational.
- 2 30 is not a perfect square. Therefore $\sqrt{30}$ has a nonterminating, nonrepeating decimal and is irrational.
- 3 Whole numbers are 0, 1, 2, 3, ... Among the choices, only 0 belongs to the whole-number set.
- 4 Let $x = 0.\overline{3}$. Then $10x = 3.\overline{3}$, so $9x = 3$ and $x = \frac{1}{3}$.
- 5 The number 45 is between 36 and 49, and it is closer to 49. Therefore $\sqrt{45}$ is closer to 7 than to 6.
- 6 Estimate $\sqrt{10} \approx 3.2$ and simplify $\sqrt{16} = 4$. The order is about 3.2, 3.5, 4.
- 7 Multiply powers with the same base by adding exponents. $3 + 8 = 11$.
- 8 Divide powers with the same base by subtracting exponents. $12 - 5 = 7$.
- 9 A power raised to another power uses multiplication of exponents. $4 \cdot 3 = 12$.
- 10 A negative exponent means reciprocal. Since $2^5 = 32$, $2^{-5} = \frac{1}{32}$.
- 11 The quotient rule gives $10^{7-3} = 10^4$. The other choices simplify to different powers.
- 12 Any nonzero number to the zero power equals 1. Therefore $8^0 = 1$, not 0.
- 13 The side length is $\sqrt{75} = 5\sqrt{3}$. Since $\sqrt{3}$ is irrational, the side length is irrational.
- 14 58 is between 49 and 64 and closer to 64, so $\sqrt{58}$ is between 7 and 8 and closer to 8. The best choice is 7.6.
- 15 First use the power rule to get $(x^3)^4 = x^{12}$. Then divide by x^5 to get $x^{12-5} = x^7$.



Get Online

Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)

- 16 Let $x = 0.1666\dots$. Then $100x = 16.666\dots$ and $10x = 1.666\dots$, so $90x = 15$ and $x = \frac{1}{6}$.

📅 Week 2 Day 4: Graphing Proportional Relationships

✔ Answers

- 1 4.5 2 2.5 3 3.5 4 Yes 5 Yes 6 No 7 $y = 4x$ 8 A 9 Yes
 10 No 11 27 12 $y = 15x$ 13 The second cyclist 14 False 15 Store B 16 10

💡 Explanations

- 1 For $y = kx$, divide y by x . $18/4 = 4.5$, so the constant of proportionality is 4.5.
- 2 For a proportional relationship, the unit rate is y/x for any point on the line with $x \neq 0$. Using $(6, 15)$, $15/6 = 2.5$.
- 3 Divide y by x for each pair. $7/2 = 3.5$, $17.5/5 = 3.5$, and $28/8 = 3.5$.
- 4 The ratio y/x is always 4. Since the constant ratio is the same, the relationship is proportional with $y = 4x$.
- 5 The graph is a straight line through the origin. That means it can be written as $y = kx$, so it represents a proportional relationship.
- 6 The graph is a line, but it does not pass through the origin. A proportional relationship must go through $(0, 0)$.
- 7 Find the constant by dividing $y/x = 20/5 = 4$. A proportional equation has the form $y = kx$, so $y = 4x$.
- 8 Only $y = 7x$ has the form $y = kx$. The other equations have added or subtracted constants or no x -term.



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



- 9 The equation is in the form $y = kx$, with $k = 0.8$. Its graph would be a straight line through the origin.
- 10 The -5 means the graph has a nonzero y -intercept. A proportional relationship must have no added or subtracted constant.
- 11 Substitute $x = 12$ into $y = 2.25x$. $2.25(12) = 27$, so $y = 27$.
- 12 The unit rate is $45/3 = 15$ pages per minute. If x is minutes and y is pages, the equation is $y = 15x$.
- 13 The first cyclist travels 14 mph. The second cyclist travels $45/3 = 15$ mph, which is faster.
- 14 A proportional relationship must pass through $(0, 0)$. If $x = 0$, then $y = k(0) = 0$, not 5.
- 15 Store A costs $18/4 = \$4.50$ per pound. Store B costs $24/6 = \$4.00$ per pound, so Store B has the lower unit price.
- 16 First find the constant of proportionality from the point: $k = 6/9 = 2/3$. Then use $y = kx$, so $y = \frac{2}{3}(15) = 10$.

📅 Week 6 Day 4: The Pythagorean Theorem

✓ Answers

- 1 15 2 25 3 12 4 12 5 17 6 24 7 Yes 8 No 9 B
- 10 False 11 No 12 12 ft 13 50 m 14 11.4 ft 15 10 16 26 cm

💡 Explanations

- 1 Use $c^2 = 9^2 + 12^2 = 81 + 144 = 225$. Since $\sqrt{225} = 15$, the hypotenuse is 15.
- 2 Use $c^2 = 7^2 + 24^2 = 49 + 576 = 625$. The square root of 625 is 25.



Get Online



Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)



- 3 Use $b^2 = 13^2 - 5^2 = 169 - 25 = 144$. The missing leg is $\sqrt{144} = 12$.
- 4 Use $b^2 = 20^2 - 16^2 = 400 - 256 = 144$. Therefore $b = 12$.
- 5 The legs are 8 and 15. $c^2 = 8^2 + 15^2 = 64 + 225 = 289$, so $c = 17$.
- 6 Subtract the known leg square from the hypotenuse square: $26^2 - 10^2 = 676 - 100 = 576$. The square root of 576 is 24.
- 7 The longest side is 26. Since $10^2 + 24^2 = 100 + 576 = 676 = 26^2$, the sides form a right triangle.
- 8 The longest side is 17. $8^2 + 14^2 = 64 + 196 = 260$, but $17^2 = 289$, so the converse is not satisfied.
- 9 Check the longest side as the hypotenuse. $6^2 + 8^2 = 36 + 64 = 100 = 10^2$, so 6, 8, 10 forms a right triangle.
- 10 The hypotenuse is opposite the right angle and is always the longest side in a right triangle.
- 11 Use the longest side 13 as the possible hypotenuse. $9^2 + 9^2 = 162$, but $13^2 = 169$, so it is not right.
- 12 The ladder is the hypotenuse. $h^2 = 15^2 - 9^2 = 225 - 81 = 144$, so $h = 12$ ft.
- 13 The diagonal is the hypotenuse of a right triangle with legs 30 and 40. $30^2 + 40^2 = 2500$, so the diagonal is 50 m.
- 14 Use $c^2 = 3^2 + 11^2 = 130$. Then $c = \sqrt{130} \approx 11.4$ ft.
- 15 The horizontal change is 6, and the vertical change is 8. The distance is $\sqrt{6^2 + 8^2} = \sqrt{100} = 10$.
- 16 First the base diagonal is $\sqrt{6^2 + 8^2} = 10$. Then the space diagonal is $\sqrt{10^2 + 24^2} = \sqrt{676} = 26$ cm.



Get Online

Find more at
[ViewMath.com/Grade8](https://www.viewmath.com/Grade8)

THANK YOU

Enjoyed This Preview?

Get the Full Book!

This preview shows just a small sample of what's inside.

The complete book includes:

- ✓ All chapters and topics
- ✓ Hundreds of practice problems
- ✓ Complete answer key with explanations
- ✓ Colorful visuals and step-by-step examples
- ✓ Reference sheets and progress trackers

 Visit us at [ViewMath.com](https://www.viewmath.com) for free resources and more books!